

A NOTE ON THE MORDELL-WEIL RANK MODULO n

TIM AND VLADIMIR DOKCHITSER

ABSTRACT. Conjecturally, the parity of the Mordell-Weil rank of an elliptic curve over a number field K is determined by its root number. The root number is a product of local root numbers, so the rank modulo 2 is (conjecturally) the sum over all places of K of a function of elliptic curves over local fields. This note shows that there can be no analogue for the rank modulo 3, 4 or 5, or for the rank itself. In fact, standard conjectures for elliptic curves imply that there is no analogue modulo n for any $n > 2$, so this is purely a parity phenomenon.

Surely, otherwise somebody would have spotted it by now.
— *Tom Fisher*

It is a consequence of the Birch–Swinnerton-Dyer conjecture that the parity of the Mordell-Weil rank of an elliptic curve E over a number field K is determined by its root number, the sign in the functional equation of the L -function. The root number is a product of local root numbers, which leads to a conjectural formula of the form

$$\mathrm{rk} E/K \equiv \sum_v \lambda(E/K_v) \pmod{2},$$

where λ is an invariant of elliptic curves over *local* fields, and v runs over the places of K . One might ask whether there is a local expression like this for the rank modulo 3 or modulo 4, or even for the rank itself. The purpose of this note is to show that, unsurprisingly, the answer is ‘no’.

The idea is simple: if the rank modulo n were a sum of local $\mathbb{Z}/n\mathbb{Z}$ -valued invariants, then $\mathrm{rk} E/K$ would be a multiple of n whenever E is defined over $\mathbb{Q}$ and $K/\mathbb{Q}$ is a Galois extension where every place of $\mathbb{Q}$ splits into a multiple of n places. However, for small $n > 2$ it is easy to find E and K for which this property fails (Theorem 2). In fact, if one believes the standard heuristics concerning ranks of elliptic curves in abelian extensions, it fails for every $n > 2$ and every $E/\mathbb{Q}$ (Theorems 9, 13).

This kind of argument can be used to test whether a global invariant has a chance of being a sum of local terms. We will apply it to other standard invariants of elliptic curves and show that the parity of the 2-Selmer rank, the parity of the rank of the p -torsion and $\dim_{\mathbb{F}_2} \mathrm{III}[2]$ modulo 4 cannot be expressed as a sum of local terms (Theorem 6). Finally, we will also comment on L -functions all of whose local factors are n^{th} powers and discuss the parity of the analytic rank for non-self-dual twists of elliptic curves (Remarks 4,7).

Our results only prohibit an expression for the rank as a *sum* of local terms. Local data does determine the rank, see Remark 15.

1. MORDELL-WEIL RANK IS NOT A SUM OF LOCAL INVARIANTS

Definition. Suppose $(K, E) \mapsto \Lambda(E/K)$ is some global invariant of elliptic curves over number fields¹. We say it is a *sum of local invariants* if

$$\Lambda(E/K) = \sum_v \lambda(E/K_v),$$

where λ is some invariant of elliptic curves over local fields, and the sum is taken over all places of K .

Implicitly, Λ and λ take values in some abelian group A , usually $\mathbb{Z}$. Moreover $\lambda(E/K_v)$ should be 0 for all but finitely many v .

Example. If the Birch–Swinnerton-Dyer conjecture holds (or if III is finite, see [4]), then the Mordell-Weil rank modulo 2 is a sum of local invariants with values in $\mathbb{Z}/2\mathbb{Z}$. Specifically, for an elliptic curve E over a local field k write $w(E/k) = \pm 1$ for its local root number, and define λ by $(-1)^{\lambda(E/k)} = w(E/k)$. Then

$$\text{rk } E/K \equiv \sum_v \lambda(E/K_v) \pmod{2}.$$

An explicit description of local root numbers can be found in [9] and [4].

Theorem 1. *The Mordell-Weil rank is not a sum of local invariants.*

This is a consequence of the following stronger statement:

Theorem 2. *For $n \in \{3, 4, 5\}$ the Mordell-Weil rank modulo n is not a sum of local invariants (with values in $\mathbb{Z}/n\mathbb{Z}$).*

Lemma 3. *Suppose $\Lambda : (\text{number fields}) \rightarrow \mathbb{Z}/n\mathbb{Z}$ satisfies $\Lambda(K) = \sum_v \lambda(K_v)$ for some function $\lambda : (\text{local fields}) \rightarrow \mathbb{Z}/n\mathbb{Z}$. Then $\Lambda(F) = 0$ whenever F/K is a Galois extension of number fields in which the number of places above each place of K is a multiple of n .*

Proof. In the local expression for $\Lambda(F)$ each local field occurs a multiple of n times. $\square$

Proof of Theorem 2. Take $E/\mathbb{Q} : y^2 = x(x+2)(x-3)$, which is 480a1 in Cremona’s notation. Writing ζ_p for a primitive p^{th} root of unity, let

$$F_n = \begin{cases} \text{the degree 9 subfield of } \mathbb{Q}(\zeta_{13}, \zeta_{103}) & \text{if } n = 3, \\ \text{the degree 25 subfield of } \mathbb{Q}(\zeta_{11}, \zeta_{241}) & \text{if } n = 5, \\ \mathbb{Q}(\sqrt{-1}, \sqrt{41}, \sqrt{73}) & \text{if } n = 4. \end{cases}$$

Because 13 and 103 are cubes modulo one another, and all other primes are unramified in F_3 , every place of $\mathbb{Q}$ splits into 3 or 9 in F_3 . Similarly F_4 and F_5 also satisfy the assumptions of Lemma 3 with $n = 4, 5$. Hence, if the Mordell-Weil rank modulo n were a sum of local invariants, it would be

¹Meaning that if $K \cong K'$ and E/K and E'/K' are isomorphic elliptic curves (identifying K with K'), then $\Lambda(E/K) = \Lambda(E'/K')$.

$0 \in \mathbb{Z}/n\mathbb{Z}$ for E/F_n . However, 2-descent shows that $\text{rk } E/F_3 = \text{rk } E/F_5 = 1$ and $\text{rk } E/F_4 = 6$ (e.g. using Magma [1], over all minimal non-trivial subfields of F_n). $\square$

Remark 4. The L -series of the curve $E = 480a1$ used in the proof over $F = F_4 = \mathbb{Q}(\sqrt{-1}, \sqrt{41}, \sqrt{73})$ is formally a 4th power, in the sense that each Euler factor is:

$$L(E/F, s) = 1 \cdot \left(\frac{1}{1-3^{-2s}}\right)^4 \left(\frac{1}{1-5^{-2s}}\right)^4 \left(\frac{1}{1+14 \cdot 7^{-2s} + 7^{2-4s}}\right)^4 \left(\frac{1}{1+6 \cdot 11^{-2s} + 11^{2-4s}}\right)^4 \cdots$$

However, it is not a 4th power of an entire function, as it vanishes to order 6 at $s = 1$. Actually, it is not even a square of an entire function: it has a simple zero at $1 + 2.1565479...i$.

In fact, by construction of F , for any $E/\mathbb{Q}$ the L -series $L(E/F, s)$ is formally a 4th power and vanishes to even order at $s = 1$ by the functional equation. Its square root has analytic continuation to a domain including $\text{Re } s > \frac{3}{2}$, $\text{Re } s < \frac{1}{2}$ and the real axis, and satisfies a functional equation $s \leftrightarrow 2 - s$, but it is not clear whether it has an arithmetic meaning.

Lemma 5. *Suppose an invariant $\Lambda \in \mathbb{Z}/2^k\mathbb{Z}$ is a sum of local invariants. Let $F = K(\sqrt{\alpha_1}, \dots, \sqrt{\alpha_m})$ be a multi-quadratic extension in which every prime of K splits into a multiple of 2^k primes of F . Then for every elliptic curve E/K ,*

$$\Lambda(E/K) + \sum_D \Lambda(E_D/K) = 0,$$

where the sum is taken over the quadratic subfields $K(\sqrt{D})$ of F/K , and E_D denotes the quadratic twist of E by D .

Proof. In the local expression for the left-hand side of the formula each local term (λ of a given elliptic curve over a given local field) occurs a multiple of 2^k times. $\square$

Theorem 6. *Each of the following is not a sum of local invariants:*

- $\dim_{\mathbb{F}_2} \text{III}(E/K)[2] \pmod{4}$,
- $\text{rk}(E/K) + \dim_{\mathbb{F}_2} \text{III}(E/K)[2] \pmod{4}$,
- $\dim_{\mathbb{F}_2} \text{Sel}_2 E/K \pmod{2}$,
- $\dim_{\mathbb{F}_p} E(K)[p] \pmod{2}$ for any prime p .

Here III is the Tate-Shafarevich group and Sel_2 is the 2-Selmer group.

Proof. The argument is similar to that of Theorem 2:

For the first two claims, apply Lemma 5 to $E : y^2 + y = x^3 - x$ (37a1) with $K = \mathbb{Q}$ and $F = \mathbb{Q}(\sqrt{-1}, \sqrt{17}, \sqrt{89})$. The quadratic twists of E by 1, -17 , -89 , $17 \cdot 89$ have rank 1, and those by -1 , 17 , 89 , $-17 \cdot 89$ have rank 0; the twist by $-17 \cdot 89$ has $|\text{III}[2]| = 4$ and the other seven have trivial $\text{III}[2]$. The sum over all twists is therefore 2 mod 4 in both cases, so they are not sums of local invariants.

For the parity of the 2-Selmer rank and of $\dim E[2]$ apply Lemma 3 to $E : y^2 + xy + y = x^3 + 4x - 6$ (14a1) with $K = \mathbb{Q}$, $F = \mathbb{Q}(\sqrt{-1}, \sqrt{17})$ and $n = 2$. The 2-torsion subgroup of E/F is of order 2 and its 2-Selmer group over F is of order 8.

Finally, for $\dim_{\mathbb{F}_p} E[p] \pmod 2$ for $p > 2$ take any elliptic curve $E/\mathbb{Q}$ with $\text{Gal}(\mathbb{Q}(E[p])/\mathbb{Q}) \cong \text{GL}_2(\mathbb{F}_p)$, e.g. $E : y^2 = x^3 - x^2 + x$ (24a4), see [10] 5.7.2. Let K be the field obtained by adjoining to $\mathbb{Q}$ the coordinates of one p -torsion point and $F = K(\sqrt{-1}, \sqrt{17})$. Because F does not contain the p^{th} roots of unity, $\dim_{\mathbb{F}_p} E(F)[p] = 1$. So, by Lemma 3 the parity of this dimension is not a sum of local invariants. $\square$

Remark 7. The functional equation expresses the parity of the analytic rank as a sum of local invariants not only for elliptic curves (or abelian varieties), but also for their twists by self-dual Artin representations. However, for the parity of the rank of non-self-dual twists there is presumably no such expression.

For example, let χ be a non-trivial Dirichlet character of $(\mathbb{Z}/7\mathbb{Z})^\times$ of order 3. Then there is no function $(k, E) \mapsto \lambda(E/k) \in \mathbb{Z}$ defined for elliptic curves over local fields k , such that for all elliptic curves $E/\mathbb{Q}$,

$$\text{ord}_{s=1} L(E, \chi, s) \equiv \sum_v \lambda(E/\mathbb{Q}_v) \pmod 2.$$

To see this, take

$$E/\mathbb{Q} : y^2 + y = x^3 + x^2 + x \text{ (19a3), } K = \mathbb{Q}, \quad F = \mathbb{Q}(\sqrt{-1}, \sqrt{17})$$

and apply Lemma 5. The twists of E, E_{-1} and E_{17} by χ have analytic rank 0, and that of E_{-17} has analytic rank 1, adding up to an odd number.

2. EXPECTATIONS

We expect the Mordell-Weil rank modulo n not to be a sum of local terms for any $n > 2$ and any class of elliptic curves. Theorems 9 and 13 below show that this is a consequence of modularity of elliptic curves, the known cases of the Birch–Swinnerton-Dyer conjecture and standard conjectures for analytic ranks of elliptic curves.

Notation. For a prime p we write Σ_p for the set of all Dirichlet characters of order p . We say that $S \subset \Sigma_p$ has density α if

$$\lim_{x \rightarrow \infty} \frac{|\{\chi : \chi \in S \mid N(\chi) < x\}|}{|\{\chi : \chi \in \Sigma_p \mid N(\chi) < x\}|} = \alpha,$$

where $N(\chi)$ denotes the conductor of χ .

Conjecture 8 (Weak form of [3] Conj. 1.2). *For $p > 2$ and every elliptic curve $E/\mathbb{Q}$, those $\chi \in \Sigma_p$ for which $L(E, \chi, 1) = 0$ have density 0 in Σ_p .*

Theorem 9. *Let $E/\mathbb{Q}$ be an elliptic curve and p an odd prime. Assuming Conjecture 8, there is no function $k \mapsto \lambda(E/k) \in \mathbb{Z}/p\mathbb{Z}$ defined for local fields k of characteristic 0, such that for every number field K ,*

$$\mathrm{rk} E/K \equiv \sum_v \lambda(E/K_v) \pmod{p}.$$

Lemma 10. *Let p be a prime number and $S \subset \Sigma_p$ a set of characters of density 0 in Σ_p . For every $d \geq 1$ there is an abelian extension $F_d/\mathbb{Q}$ with Galois group $G \cong \mathbb{F}_p^d$, such that no characters of G are in S .*

Proof. Without loss of generality, we may assume that if $\chi \in S$ then $\chi^n \in S$ for $1 \leq n < p$. When $d = 1$, take F_1 to be the kernel of any $\chi \in \Sigma_p \setminus S$. Now proceed by induction, supposing that F_{d-1} is constructed. Writing Ψ for the set of characters of $\mathrm{Gal}(F_{d-1}/\mathbb{Q})$, the set

$$S_d = \bigcup_{\psi \in \Psi} \{\phi\psi : \phi \in S\}$$

still has density 0. Pick any $\chi \in \Sigma_p \setminus S_d$, and set F_d to be the compositum of F_{d-1} and the degree p extension of $\mathbb{Q}$ cut out by χ . It is easy to check that no character of $\mathrm{Gal}(F_d/\mathbb{Q})$ lies in S . $\square$

Proof of Theorem 9. Pick a quadratic field $\mathbb{Q}(\sqrt{D})$ such that the quadratic twist E_D of E by D has analytic rank 1, which is possible by [2, 8, 11]. By Conjecture 8, the set S of Dirichlet characters χ of order p such that $L(E_D, \chi, 1) = 0$ has density 0. Apply Lemma 10 to S with $d = 3$. Every place of $\mathbb{Q}$ splits in the resulting field $F = F_3$ into a multiple of p places ($\mathbb{Q}_l$ has no $\mathbb{F}_p^3$ -extensions, so every prime has to split).

Arguing by contradiction, suppose the rank of $E \bmod p$ is a sum of local invariants. Because in $F/\mathbb{Q}$ and therefore also in $F(\sqrt{D})/\mathbb{Q}$ every place splits into a multiple of p places,

$$\mathrm{rk} E/F \equiv 0 \pmod{p} \quad \text{and} \quad \mathrm{rk} E/F(\sqrt{D}) \equiv 0 \pmod{p}$$

by Lemma 3. Therefore $\mathrm{rk} E_D/F = \mathrm{rk} E/F(\sqrt{D}) - \mathrm{rk} E/F$ is a multiple of p . On the other hand,

$$L(E_D/F, s) = \prod_{\chi} L(E_D, \chi, s),$$

the product taken over the characters of $\mathrm{Gal}(F/\mathbb{Q})$. By construction, it has a simple zero at $s = 1$. Because F is totally real of odd degree over $\mathbb{Q}$, by Zhang's theorem [12] Thm. A, E_D/F has Mordell-Weil rank $1 \not\equiv 0 \pmod{p}$, a contradiction. $\square$

Conjecture 11 (Goldfeld [6]). *For every elliptic curve $E/\mathbb{Q}$, those $\chi \in \Sigma_2$ for which $\mathrm{ord}_{s=1} L(E, \chi, s) > 1$ have density 0 in Σ_2 .*

Conjecture 12. *Every elliptic curve $E/\mathbb{Q}$ has a quadratic twist of Mordell-Weil rank 2.*

Theorem 13. *Let $E/\mathbb{Q}$ be an elliptic curve. Assuming Conjectures 11 and 12, there is no function $k \mapsto \lambda(E/k) \in \mathbb{Z}/4\mathbb{Z}$ defined for local fields k of characteristic 0, such that for every number field K ,*

$$\mathrm{rk} E/K \equiv \sum_v \lambda(E/K_v) \pmod{4}.$$

Proof. Let $\mathbb{Q}(\sqrt{D})$ be a quadratic field such that the quadratic twist $E' = E_D$ of E by D has Mordell-Weil rank 2 (Conjecture 12). The set S of those $\chi \in \Sigma_2$ for which $\mathrm{ord}_{s=1} L(E', \chi, s) > 1$ has density 0 (Conjecture 11). Let P be the set of primes where E' has bad reduction union $\{\infty\}$, and apply Lemma 10 to S with $d = 5 + 3|P|$. The resulting field F_d has a subfield F of degree 2^5 over $\mathbb{Q}$, where all places in P split completely: $\mathbb{Q}_l$ has no $\mathbb{F}_2^4$ -extensions, so the condition that a given place in P splits completely drops the dimension by at most 3. By the same argument, every place of $\mathbb{Q}$ splits in F into a multiple of 4 places.

Arguing by contradiction, suppose the rank of $E \bmod 4$ is a sum of local invariants. Because in $F/\mathbb{Q}$ and therefore also in $F(\sqrt{D})/\mathbb{Q}$ every place splits into a multiple of 4 places,

$$\mathrm{rk} E/F \equiv 0 \pmod{4} \quad \text{and} \quad \mathrm{rk} E/F(\sqrt{D}) \equiv 0 \pmod{4}$$

by Lemma 3. Therefore $\mathrm{rk} E'/F = \mathrm{rk} E/F(\sqrt{D}) - \mathrm{rk} E/F$ is a multiple of 4. Now we claim that E'/F has rank 2 or 33, yielding a contradiction.

Let $\mathbb{Q}(\sqrt{m}) \subset F$ be a quadratic subfield. The root number of E' over $\mathbb{Q}(\sqrt{m})$ is 1, because the root number is a product of local root numbers and the places in P split in $\mathbb{Q}(\sqrt{m})$. (The local root number is +1 at primes of good reduction.) So

$$L(E'/\mathbb{Q}(\sqrt{m}), s) = L(E'/\mathbb{Q}, s)L(E'_m/\mathbb{Q}, s)$$

vanishes to even order at $s = 1$. Hence the 31 twists of E' by the non-trivial characters of $\mathrm{Gal}(F/\mathbb{Q})$ have the same analytic rank 0 or 1, by the choice of F . By Kolyvagin's theorem [7], their Mordell-Weil ranks are the same as their analytic ranks, and so $\mathrm{rk} E'/F$ is either $2 + 0$ or $2 + 31$. $\square$

Remark 14. In some cases, it may seem reasonable to try and write some global invariant in $\mathbb{Z}/n\mathbb{Z}$ as a sum of local invariants in $\frac{1}{m}\mathbb{Z}/n\mathbb{Z}$, i.e. to allow denominators in the local terms. For instance, one could ask whether the parity of the rank of a cubic twist (as in Remark 7) can be written as a sum of local invariants of the form $\frac{a}{3} \bmod 2\mathbb{Z}$.

However, introducing a denominator does not appear to help. First, the prime-to- n part m' of m adds no flexibility, as can be seen by multiplying the formula by m' . (For instance, if there were a formula for the parity of the rank of a cubic twist as a sum of local terms in $\frac{a}{3} \bmod 2\mathbb{Z}$, then multiplying it by 3 would yield a formula for the same parity with local terms in $\mathbb{Z}/2\mathbb{Z}$.) As for the non-prime-to- n part, e.g. the proofs of Theorems 9 and 13 immediately adapt to local invariants in $\frac{1}{p^k}\mathbb{Z}/p\mathbb{Z}$ and $\frac{1}{2^k}\mathbb{Z}/4\mathbb{Z}$, by increasing d by k .

Remark 15. The negative results in this paper rely essentially on the fact that we allow only additive formulae for global invariants in terms of local invariants. Although Theorem 1 shows that there is no formula of the form

$$\mathrm{rk} E/K = \sum_v \lambda(E/K_v),$$

the Mordell-Weil rank *is* determined by the set $\{E/K_v\}_v$ of curves over local fields. In other words,

$$\mathrm{rk} E/K = \text{function}(\{E/K_v\}_v).$$

In fact, for any abelian variety A/K the set $\{A/K_v\}_v$ determines the L -function $L(A/K, s)$ which is the same as $L(W/\mathbb{Q}, s)$ where W is the Weil restriction of A to $\mathbb{Q}$. By Faltings' theorem [5] the L -function recovers W up to isogeny, and hence also recovers the rank $\mathrm{rk} A/K (= \mathrm{rk} W/\mathbb{Q})$.

Acknowledgements. The first author is supported by a Royal Society University Research Fellowship. The second author would like to thank Gonville & Caius College, Cambridge.

REFERENCES

- [1] W. Bosma, J. Cannon, C. Playoust, The Magma algebra system. I: The user language, *J. Symb. Comput.* 24, No. 3–4 (1997), 235–265.
- [2] D. Bump, S. Friedberg, and J. Hoffstein, Nonvanishing theorems for L -functions of modular forms and their derivatives, *Invent. Math.* 102 (1990), 543–618.
- [3] C. David, J. Fearnley, H. Kisilevsky, Vanishing of L -functions of elliptic curves over number fields, in *Ranks of Elliptic Curves and Random Matrix Theory*, LMS Lecture Note Series, 341, Cambridge University Press (2007), 247–259.
- [4] T. Dokchitser, V. Dokchitser, Root numbers and parity of ranks of elliptic curves, 2009, arxiv: 0906.1815.
- [5] G. Faltings, Endlichkeitssätze für abelsche Varietäten über Zahlkörpern, *Invent. Math.* 73 (1983), no. 3, 349–366.
- [6] D. Goldfeld, Conjectures on elliptic curves over quadratic fields, *Springer Lect. Notes* 751 (1979), 108–118.
- [7] V. A. Kolyvagin, Euler systems, *The Grothendieck Festschrift*, *Prog. in Math.*, Boston, Birkhauser (1990).
- [8] M. R. Murty and V. K. Murty, Mean values of derivatives of modular L -series, *Annals of Math.* 133 (1991), 447–475.
- [9] D. Rohrlich, Galois Theory, elliptic curves, and root numbers, *Compos. Math.* 100 (1996), 311–349.
- [10] J.-P. Serre, Propriétés galoisiennes des points d'ordre fini des courbes elliptiques, *Invent. Math.* 15 (1972), no. 4, 259–331.
- [11] J.-L. Waldspurger, Correspondences de Shimura et quaternions, *Forum Math.* 3 (1991), 219–307.
- [12] S. Zhang, Heights of Heegner points on Shimura curves, *Annals of Math.* 153 (2001), 27–147.

ROBINSON COLLEGE, CAMBRIDGE CB3 9AN, UNITED KINGDOM
E-mail address: t.dokchitser@dpms.cam.ac.uk

EMMANUEL COLLEGE, CAMBRIDGE CB2 3AP, UNITED KINGDOM
E-mail address: v.dokchitser@dpms.cam.ac.uk